

## SURGICAL APPROACH AND OUTCOMES IN SALIVARY GLAND SWELLING

Dr. Akshay Akulwar<sup>1</sup>, Dr. Akshay Bavikatte<sup>2</sup>

<sup>1</sup>Assistant Professor, Mahatma Gandhi Institute of Medical Sciences Sewagram Wardha, Maharashtra.

<sup>2</sup>P.G. J.J.M. Medical College Davangere Karnataka.

**Article Info:** Received 10 January 2020; Accepted 01 February 2020

**DOI:** <https://doi.org/10.32553/ijmbs.v4i2.1030>

**Corresponding author:** Dr. Akshay Bavikatte

**Conflict of interest:** No conflict of interest.

### Abstract

**Introduction:** Salivary gland tumours are rare tumours and majority of these are benign and about 20% are malignant. The incidence of salivary gland cancers ranges from 0.5 to 2 per 100,000 in different parts of the world. Salivary gland swellings in the parotid or submandibular glands usually present as an enlarging mass and may be associated with neurological symptoms like facial nerve paralysis or pain. Minor salivary gland tumours present as a submucosal intraoral mass which subsequently ulcerates. Acute inflammatory conditions generally can be diagnosed by history and physical examination alone, whereas chronic inflammatory diseases and granulomatous disorders require supplemental diagnostic information including lab tests, imaging studies and biopsy. FNAC of salivary gland tumours is advantageous to both the patient and the clinician because of its immediate results, accuracy, lack of complications and economy. Appropriate therapeutic management may be planned earlier, whether it is local excision for a benign neoplasm, radical surgery for a malignant one or any other alternate treatment.

**Material and Methods:** All patients admitted to surgical wards due to obstructions of the salivary duct and neoplasia were included in the study. Demographic data from all the patients were collected, also they were evaluated for clinical examination, routine laboratory investigations and specific investigations. After evaluation of the swellings by clinical examination and by specific investigations, a surgical plan was formulated. The final decision was taken by the operative surgeon.

**Results:** 40 cases of salivary gland swellings were admitted during study period. Age of the patients varied from 9 years to 80 years. 62.5% (25 cases) were found in the parotid gland, 30% cases (12) in submandibular gland and 7.5% cases (3) in the sublingual gland. Surgery was the treatment for all cases of tumors. Superficial parotidectomy was done in all the 21 cases of parotid tumour (56.41%) without deep lobe involvement and total parotidectomy was done in 3 cases (7.6%) with deep lobe involvement. In all the cases of submandibular gland lesions, excision of submandibular gland was done. Excision of the sublingual gland was done in 3 cases of ranula. One case of adenoid cystic carcinoma was referred to higher center because of the advanced malignancy.

**Conclusion:** Salivary gland tumors occur more commonly in the parotid gland, most often benign, pleomorphic adenoma constitute majority of all neoplasm. Surgery is the main modality of treatment in salivary gland sialolithiasis. Most commonly done surgery is excision of submandibular salivary gland & superficial parotidectomy and also for salivary gland tumors. Awareness and early referral is necessary, as prognosis is good if treated early.

**Keywords:** SALIVARY GLAND, TUMOUR, NEOPLASM, SURGERY

### Introduction

Salivary gland tumours are rare tumours and majority of these are benign and about 20% are malignant. The incidence of salivary gland cancers ranges from 0.5 to 2 per 100,000 in different parts of the world, with the highest incidence seen in Croatia<sup>i</sup>. These tumours can occur in both the major and minor salivary glands. About 80% of major salivary gland tumours are seen in the parotid glands, while most minor salivary tumours are located in the palate<sup>ii</sup>. 20–25% of the parotid glands tumours are malignant. Prevalence is about 40% for the submandibular glands, and more than 90% of sublingual gland tumours are malignant<sup>iii iv</sup>.

Some authors concluded that a diet rich in vitamin C and low in cholesterol may be effective in preventing salivary gland cancer<sup>v</sup>. Immunosuppression, radiation and HIV infection was seen to increase the risk of salivary gland cancers<sup>vi vii</sup>.

Salivary gland swellings in the parotid or submandibular glands usually present as an enlarging mass and may be associated with neurological symptoms like facial nerve paralysis or pain. Minor salivary gland tumours present as a submucosal intraoral mass which subsequently ulcerates<sup>viii</sup>.

Acute inflammatory conditions generally can be diagnosed by history and physical examination alone, whereas chronic inflammatory diseases and granulomatous disorders require supplemental diagnostic information including lab tests, imaging studies and biopsy. Accurate pathological diagnosis is necessary for proper management of neoplastic disorders<sup>ix</sup>. These tumours usually occur in adults with a female predominance, but about 5% occur in children less than 16years<sup>x</sup>.

FNAC of salivary gland tumours is advantageous to both the patient and the clinician because of its immediate results, accuracy, lack of complications and economy<sup>9</sup>. Appropriate therapeutic management may be planned earlier, whether it is local excision for a benign neoplasm, radical surgery for a malignant one or any other alternate treatment. In general, disregarding specific histological types, Prognosis is most favorable in those located in palate, less favorable in parotid and least favorable in submandibular gland. With non-neoplastic lesions, metastasis and lymph proliferative disorders, conservative management, chemotherapy or radiotherapy might be respectively preferable<sup>xi</sup>

With this background present study was planned to study the methods of current surgical treatment and outcomes of salivary glands swellings.

#### Material and Methods:

The present prospective, observational study was carried out in Department of Surgery from May 2009 to July 2011 in J.J.M. Medical College and Chigateri District hospital, Davangere.

All patients admitted to surgical wards of J.J.M. Medical College with salivary gland swellings due to obstructions of the salivary duct and neoplasia were included in the study. Patients were excluded if salivary gland swellings arising as a result of congenital conditions, salivary swellings associated with systemic diseases and salivary gland swellings arising as a result of inflammation like Mumps or Parotitis.

Demographic data from all the patients were collected, also they were evaluated for clinical examination, routine laboratory investigations and specific investigations. In history, importance was given to presenting complaints, duration of lump, rapid increase in size, associated symptoms of facial nerve involvement, previous surgical. Associated medical conditions like diabetes, hypertension and anemia were managed and controlled before surgery with the patient's advice.

All routine investigations were carried out on all the patients. After evaluation of the swellings by clinical

examination and by specific investigations, a surgical plan was formulated. The final decision was taken by the operative surgeon. The required specimen was sent for histopathological examinations. Appropriate antibiotics and analgesics are administered post operatively for all cases. Drainage tube was removed when the drain was less than 20ml and sutures were removed on 5<sup>th</sup> day. Malignant tumors were referred to Kidwai Memorial Institute of Oncology, after surgery, for postoperative radiotherapy. The adjuvant treatment was decided depending on the final HPE report. Different modalities of treatment adopted in this study are either surgery or Surgery and post-operative radiotherapy.

The follow up period of these patients ranged from 3months to 1 year. All patients were asked for follow up after 15 days of surgery then every month for 1<sup>st</sup> year then every 3 months in 2<sup>nd</sup> year, to detect morbidity and recurrence.

All data was entered in the Excel sheet and master chart was prepared. Statistical analysis was done by SPSS version 12.0.

#### Results:

40 cases of salivary gland swellings were admitted during study period. Age of the patients varied from 9 years to 80 years. Average age of the patient was 40.6 years. The case of lowest age group i.e., 9 years was observed with non-inflammatory swelling and the case of highest age i.e., 80 years was of tumor swelling. Out of 40 cases 15(35%) cases were male and 25(65%) cases were female.

**Table 1: Mode of Clinical Presentation**

Mode	No. of Cases	Percentage
Swelling	40	100.0
Pain	26	65.0
Fever	8	20.0
Increased salivation	11	27.5
Tenderness	22	55.0
Fixity of swelling	4	10.0
Ear lobe elevation	19	47.5
Deep lobe involvement	3	7.5
Facial nerve paralysis	1	2.8

All cases presented with, symptoms of swelling (100%), 65 % (26) presented with pain. 55 % (22) presented with tenderness. Three cases were with deep lobe involvement (11.4%), 19 cases of ear lobe elevation (47.5%). Facial nerve paralysis occurred in one case (2.8%).

**Table 2: Site for Various Salivary Gland Swellings**

No. of cases	Parotid	Submandibular	Sublingual
40	25 (62.5%)	12(30.0%)	3(7.5%)

In our study, 62.5% (25 cases) were found in the parotid gland, 30% cases (12) in submandibular gland and 7.5% cases (3) in the sublingual gland.

In our study, 41.66% (5) of cases of sialolithiasis were in right submandibular gland, 58.33% (7) of cases in the left submandibular gland and 3 cases (100%) of ranula were seen in right sublingual gland only. Out of 25 salivary tumors, 96.1% were benign and 2.5% malignant. Out of 25 cases of parotid tumors, 22 (88.46%) cases were seen in superficial and 3 (11.53%) in deep lobe

**Table 3:** Incidence of Superficial and Deep Lobe Involvement of Parotid Gland Tumours

No of Tumors	Superficial Lobe	Deep Lobe
25	22(88.46%)	3(11.53%)

**Table 4:** Correlation of FNAC and Histopathology

Lesions	No. of Patients	FNAC (%)	BIOPSY (%)
Pleomorphic adenoma	22	100	100
Warthin tumour	3	100	100
Adenoid cystic Ca.	1	-	-

The accuracy of FNAC was 100% in case of benign salivary gland tumours. One case which was diagnosed by FNAC as adenoid cystic carcinoma was referred to higher center for the management.

**Table 5:** Surgical Procedures Adopted for Various Salivary Gland Swellings

Procedures	No. of Patients	Percentage
Excision of submandibular gland	12	28.20
Superficial parotidectomy	21s	56.41
Total Parotidectomy	3	7.6
Excision ranula	3	7.6
Total	39	100

Surgery was the treatment for all cases of tumors. Superficial parotidectomy was done in all the 21 cases of parotid tumour (56.41%) without deep lobe involvement and total parotidectomy was done in 3 cases (7.6%) with deep lobe involvement. In all the cases of submandibular gland lesions, excision of submandibular gland was done. Excision of the sublingual gland was done in 3 cases of ranula. One case of adenoid cystic carcinoma was referred to higher center because of the advanced malignancy.

**Table 6:** Post Operative Complications

Nature of Complications	No. of Patients	Percentage
Facial nerve paralysis	1	2.5
Mandibular nerve paralysis	1	2.5
Wound infection	8	20

Post operative complications in my study of 40 cases were low. One case of facial nerve paralysis occurred after parotid tumour surgery in the case of deep lobe

involvement and one case of mandibular nerve palsy occurred with submandibular sialadenectomy, wound infection was noticed in 8 cases.

### Discussion:

History of salivary gland disease date backs to times of Hippocrates. Although parotid gland has been surgically approached on selective basis for at least the last 300years, an understanding of parotid anatomy, especially in relation to the facial nerve, was not made clear until early part of 20 century. Earliest reports of parotid extirpative surgery were recorded in Dutch literature of late 1600<sup>xii</sup>. According to Foote and Frazel, term mixed tumour dates from Minssen's review in 1874- Which is cited by Ahlbom. This neoplasm was originally designated the benign mixed tumour in 1866. A name change to pleomorphic adenoma was suggested in 1948<sup>xiii</sup>.

Out of 40 cases in our study 15(35%) cases were male and 25(65%) cases were female. This finding was in accordance with the study by Ansari<sup>5</sup> in Iran, Otoh *et al*<sup>xiv</sup> in Nigeria. In our study, incidence percentage of sialolithiasis i.e., 12 cases were found in submandibular gland which co-relates with Pizzirani *et al*<sup>xv</sup> in 1985 and J. Lustmann *et al*<sup>xvi</sup> in 1990.

In the present study 97.5% tumours were benign and 2.5% were malignant. Renehan *et al*<sup>xvii</sup> 1996 observed 80% benign and 0% malignant tumour in his study while Skolnik *et al*<sup>xviii</sup> in 1977 observed 59.40% benign and 30.60% malignant tumour. In our study, all the salivary gland tumors were observed in parotid gland. Comparative study was in accordance to Renehan *et al*<sup>17</sup>.

**Table 7:** Frequency of Benign and Malignant Salivary Tumours In Different Series

Series	No. of tumors	Benign	Malignant
Foote <i>et al</i> <sup>xix</sup> 1954	730	68.30%	31.70%
Skolnik <i>et al</i> <sup>xx</sup> 1977	435	59.40%	30.60%
Khazanchi <i>et al</i> <sup>xxi</sup> 1988	88	63.60%	36.40%
Renehan <i>et al</i> <sup>17</sup> 1996	1194	80.00%	0.00%
Present study	25	97.5%	2.5%

In accordance with the observation in other series, the benign tumors predominate in our study.

**In this study**, out of 25 parotid tumours, 22 (88.46%) were seen in superficial lobe of parotid and 3 (11.53%) in deep lobe. In a study by H. Leverstein *et al*<sup>xxii</sup>, superficial tumours were observed in 192(78.3%) cases and deep tumours in 54(22%) cases.

In our study of 40 cases of salivary gland swelling, shows that, surgery is the treatment of choice in all cases of salivary gland swellings. FNAC plays an important role in the diagnosis of salivary gland tumors and accuracy rate was 100% in our series. Benign swelling of the salivary gland found in lower decade of life, whereas, malignant swelling was found in 8th decade of life

### Conclusion:

Salivary gland tumors occur more commonly in the parotid gland, most often benign, pleomorphic adenoma constitute majority of all neoplasm. History and physical examination complement FNAC and help in diagnosis. FNAC has good accuracy in diagnosing salivary gland swellings. Surgery is the main modality of treatment in salivary gland sialolithiasis. Most commonly done surgery is excision of submandibular salivary gland & superficial parotidectomy and also for salivary gland tumors. Since most malignant tumors is asymptomatic and long standing benign tumors can undergo malignant change, community awareness and early referral is necessary, as prognosis is good if treated early

### References:

1. Fifty years of cancer incidence: CI5 I-IX. *Int J Cancer*. 2010 Dec 15; 127(12):2918-27. Parkin DM, Ferlay J, Curado MP, Bray F, Edwards B, Shin HR, Forman D
2. Speight PM, Barrett AW. Salivary gland tumours. *Oral Dis*. 2002 Sep; 8(5):229-40.
3. Arshad AR. Parotid swellings: report of 110 consecutive cases. *Med J Malaysia*. 1998 Dec; 53(4):417-22.
4. Loyola AM, de Araújo VC, de Sousa SO, de Araújo NS. Minor salivary gland tumours. A retrospective study of 164 cases in a Brazilian population. *Eur J Cancer B Oral Oncol*. 1995 May; 31B(3):197-201.
5. Horn-Ross PL, Morrow M, Ljung BM. Diet and the risk of salivary gland cancer. *Am J Epidemiol*. 1997 Jul 15; 146(2):171-6.
6. Sun EC, Curtis R, Melbye M, Goedert JJ. Salivary gland cancer in the United States. *Cancer Epidemiol Biomarkers Prev*. 1999 Dec; 8(12):1095-100.
7. Dong C, Hemminki K. Second primary neoplasms among 53, 159 haematolymphoproliferative malignancy patients in Sweden, 1958-1996: a search for common mechanisms. *Br J Cancer*. 2001 Sep 28; 85(7):997-1005.
8. To VS, Chan JY, Tsang RK, Wei WI. Review of salivary gland neoplasms. *ISRN Otolaryngol*. 2012;2012:872982.
9. Cummings CW. "Otolaryngology-Head and neck surgery". Chapter 56-63: 2nd edition, Vol. 2, Mosby, 1993
10. Young JA. Salivary glands. Chapter 6. In: *Fine needle Aspiration Cytopathology*. Jennifer A Young. 1 edition. London. Blackwell Scientific Publications; 1993: pp.48-67.
11. Cohen MB, Fisher PE, Holly EA, Ljung BM, Lowhagen T, Bottles K. Fine needle aspiration biopsy diagnosis of Mucoepidermoid Carcinoma-statistical analysis. *Acta Cytol* 1990; 1 (34): 43-49
12. Rankow RM, Polayer IM. Disease of the salivary glands Philadelphia. W B Saunders. 1996; 1-16.
13. Spiro RH, Huros AG and Strong EW. Cancer of the parotid gland. A Clinicopathological study of 288 primary cases. *Am J Surg* 1975; 30; 452- 459.
14. Otoh EC, Johnson NW, Olasoji H, Danfillo IS, Adeleke OA. Salivary gland neoplasms in Maiduguri, north-eastern Nigeria. *Oral Dis*. 2005 Nov; 11(6):386-91.
15. Pizzirani C et al. Considerazioni, clinico statistiche SU 102 Casi Di Calcolosi Salivare. *Mondo Odontostomatol* 1985; 27: 41-51
16. Lustmann J et al. A survey on 245 patients and a review of the literature. *Int J Oral Maxillofac Surg* 1990; 19: 135-138s
17. Renehan A et al. Long term follow up of over 1000 patients with salivary gland tumours treated in single centre. *Br J Surg* 1996; 83: 1750-1754
18. Skolnik EM. Tumours of major salivary glands. *Laryngoscope* 1977; 84: 843- 61.
19. Foote JW et al. Tumours of major salivary glands. In *Atlas of tumours pathology*. Sec. 4, Fascicle 11. Washington DC, 1954, Armed forces institute of pathology
20. Skolnik EM. Tumours of major salivary glands. *Laryngoscope* 1977; 84: 843- 61
21. Khanzanchi RK et al. Tumours of parotid gland: A review of 88 patients and current methods of treatment. *Ind J Cancer* 1988; 25 (1): 1-6
22. Leverstein H et al. Surgical management of 246 previously untreated pleomorphic adenomas of the parotid gland. *Br J Surg* 1997; 84(3): 399-403