

TO STUDY THE THERAPEUTIC EFFICACY OF THE GLYCOLIC ACID PEELING, SALICYLIC ACID PEELING AND MICRO- DERMABRASION IN THE TREATMENT OF MELASMA

Dr. Manish Kumar Meena¹, Dr. Deepika Meena²

^{1,2} Senior Resident

¹Department of Dermatology, Sardar Patel Medical College & Associate Group of PBM Hospitals, Bikaner (Rajasthan)

²Department of Dental, Sardar Patel Medical College & Associate Group of PBM Hospitals, Bikaner (Rajasthan)

Article Info: Received 10 May 2020; Accepted 20 June 2020

DOI: <https://doi.org/10.32553/ijmbs.v4i6.1225>

Corresponding author: Dr. Deepika Meena

Conflict of interest: No conflict of interest.

Abstract

Background: This study is aimed the therapeutic efficacy of the glycolic acid peeling, salicylic acid peeling and micro-dermabrasion in the treatment of melasma.

Methods: The present study was carried out in the department of dermatology in a teaching institute. A total of 90 patients were enrolled for the study over a period of one year.

Results: In those who were treated with GA peeling; 20.00 % had excellent response, 30.00 % had good response, 36.67% had fair response and only 13.33% had poor response. Out of total 30 patients treated with SA peeling 16.67% had excellent response, 20.00% had good response, 40.00% had fair response and 23.33% had poor response. In this group no patient had excellent response while good response was seen in 26.67%, fair response in 36.67% and poor response in another 36.67% patients.

Conclusion: all three agents were effective and safe in Indian patients, with Glycolic Acid Peel being better effect on melasma patients.

Keywords: *Therapeutic, Melasma, Pregnancy.*

Introduction

It is a chronic and recurrent disorder which has been under-diagnosed and under-treated due to lack of effective therapies and the perception that it is merely a cosmetic nuisance. The main area of involvement is face and hence it is of major cosmetic concern to the patient.

The exact prevalence of melasma is unknown in most of the countries. Melasma is a very common cutaneous disorder, accounting for 0.25 to 4% of the patients seen in Dermatology Clinics in South East Asia, and is the most common pigment disorder among Indians¹.

Melasma is primarily a disease of women of child-bearing age although 10% of cases occur in men. Many physicians outside the dermatological community have not been aware of available treatments and therefore, have typically not addressed the problem unless asked or have reassured their patients that the hyper-pigmentation would fade after delivery.

The exact causes of melasma are unknown. However, multiple factors are implicated in its etiopathogenesis, mainly sunlight, genetic predisposition, and role of female hormonal activity. Exacerbation of melasma is almost

inevitably seen after uncontrolled sun exposure and conversely melasma gradually fades during a period of sun avoidance. Genetic factors are also involved, as suggested by familial occurrence and the higher prevalence of the disease among Hispanics and Asians².

This study is aimed the therapeutic efficacy of the glycolic acid peeling, salicylic acid peeling and micro-dermabrasion in the treatment of melasma.

Methodology

Patient selection criteria: Patients of all types of melasma attending the outpatient department of dermatology were selected irrespective of age, sex, duration and previous therapy.

Patient exclusion criteria: Certain parameters were binding while selecting the patients for the study which were strictly taken care of, during the enrollment procedure.

- Pregnancy and lactating mothers
- Past history of Herpes Labialis and Herpes Zoster
- Photosensitivity
- Known keloidal tendency in the patient
- Active acne lesions over face

- History of sensitivity to Al₂O₃
- History of taking oral isotretinoin in last 1 year
- Old facial dyschromia
- Ashy melanosis
- Uncontrolled systemic disease
- Any condition necessitating UV -light therapy
- Any concomitant disease that might interfere with the diagnosis of facial hyper-pigmentation.

Study comprised of total 60 patients who satisfactorily fulfilled the inclusion and exclusion criteria.

Method of Data Collection

- Total 90 patients were enrolled for the study.
- Informed consent was taken and counseling was done with proper explanation of the whole procedure, effects and probable side effects, before starting the procedure.
- Detailed history was taken and proper examination was done.
- Each patient was offered one of the following three regimens:
 - Regimen I - 30% GA
 - Regimen II - 20% SA
 - Regimen III - Micro-dermabrasion
- Patients following regimen 1 & 2 were primed with 0.025% tretinoin cream applied at night time for 10 minutes in first 15 days.

All patients were prescribed a sunscreen lotion > SPF -30 and its importance was explained for both during and after the treatment.

Results

The present study comprises of 90 patients of melasma. Out of these 30 patients were treated with GA peel, 30 with SA peel and the remaining 30 were treated with micro-dermabrasion. Patients were evaluated and analyzed after 6 sittings [3 months] each of which was carried out 2 weeks apart.

Table 1: Response with Glycolic Acid Peel:

Response	Total	Percentage
Excellent	6	20.00
Good	9	30.00
Fair	11	36.67
Poor	4	13.33

In those who were treated with GA peeling; 20.00 % had excellent response, 30.00 % had good response, 36.67% had fair response and only 13.33% had poor response.

Table 2: Response with Salicylic Acid Peel:

	Total	Percentage [%]
Excellent	5	16.67
Good	6	20.00
Fair	12	40.00
Poor	7	23.33

Out of total 30 patients treated with SA peeling 16.67% had excellent response, 20.00% had good response, 40.00% had fair response and 23.33% had poor response.

Table 3: Response With Micro-Dermabrasion:

Response	Total	Percentage [%]
Excellent	0	0
Good	8	26.67
Fair	11	36.67
Poor	11	36.67

In this group no patient had excellent response while good response was seen in 26.67%, fair response in 36.67% and poor response in another 36.67% patients.

Discussion

In those who were treated with GA peeling; 20.00 % had excellent response, 30.00 % had good response, 36.67% had fair response and only 13.33% had poor response. A lot of studies are carried out previously on efficacy of GA peeling in melasma.

A similar study was carried out on 10 Asian women by Lim J.T. and Tham S.N.³. Statistically significant improvement was noted in 40% of the patients.

Javaheri et al⁴ performed a study with 25 Indian women with melasma. Overall improvement in melasma (reduction in MASI) was observed in 91% patients.

In the present study patients with epidermal type of melasma demonstrated a better response to GA peel treatment than those with mixed type/dermal type melasma which is in compliance to the study by Javaheri et al⁴.

In our study out of total 20 patients treated with SA peeling 16.67% had excellent response, 20.00% had good response, 40.00% had fair response and 23.33% had poor response.

Grimes⁵ conducted a similar study on 6 patients of melasma with SA peel and there was improvement in 66%.

In response with micro-dermabrasion no patient had excellent response while good response was seen in 26.67%, fair response in 36.67% and poor response in another 36.67% patients

Kunachak, Leelaudomlipi, and Wongwaisayawan⁶ have reported a large-scale study on dermabrasion in 533 patients with facial melasma. One study was conducted by Alam Murad and Omura Nayomi⁷ comparing GA peels with micro-dermabrasion. Result of my study shows similar response with both modalities.

Conclusion

Females were affected more commonly during their late third decade of life. Although we did not find the exact cause of melasma, we noticed that sun-exposure, pregnancy, and taking of oral contraceptive pills could precipitate or exacerbate the melasma.

Bibliography

1. Bandyopadhyay D. Topical treatment of melasma. *Indian J Dermatol.* 2009;54:303–92.
2. Pasricha JS, Khaitan BK, Dash S. Pigmentary disorders in India. *Dermatol Clin.* 2007;25:343–522.
3. Estrada-Castanon R, Torres-Bibiano B, Alarcon-Hernandez H. Epidemiologia cutanea en dos sectores de atencion medica en Guerrero, Mexico. *Dermatologia Rev Mex.* 1992;36:29–34.
4. Failmezger C. Incidence of skin disease in Cazco, Peru. *Int J Dermatol.* 1992;36:29–34.
5. Goh CL, Dlova CN. A retrospective Study on the clinical presentation and treatment outcome of melasma in a tertiary dermatological referral centre in Singapore. *Singapore Med J.* 1999;40:455–8.
6. Katsambas AD, Stratigos AJ, Lotti TM. Melasma. In: Katsambas AD, Lotti TM, editors. *European handbook of dermatological treatments.* 2nd ed. Berlin: Springer; 2003. pp. 336–41.
7. Resnik S. Melasma induced by oral contraceptive drug. *JAMA.* 1967;199:601–5.
8. 15. Vazquez M, Maldonado H, Benmaman C. Melasma in men, a clinical and histological study. *Int J Dermatol.* 1988;27:25–7.
9. 16. Sanchez NP, Pathak MA, Sato S, Fitzpatrick TB, Sanchez JL, Mihm MC., Jr Melasma: A clinical, light microscopic, ultrastructural and immunofluorescence study. *J Am Acad Dermatol.* 1981;4:698–710.
10. 17. Pathak MA. Clinical and therapeutic aspects of Melasma: An overview. In: Fitz Patrick TB, Wick MM, Toda K, editors. *Brown melanoderma.* Tokyo: University of Tokyo Press; 1986. pp. 161–72.
11. Katsambas A, Antoniou C. Melasma: Classification and treatment. *J Eur Acad Dermatol Venereol.* 1995;4:217–23.
12. Thappa DM. Melasma (chloasma): A review with current treatment options. *Indian J Dermatol.* 2004;49:165–76.
13. Goh CL, Dlova CN. A retrospective Study on the clinical presentation and treatment outcome of melasma in a tertiary dermatological referral centre in Singapore. *Singapore Med J.* 1999;40:455–8.
14. Nicolaidou E, Antoniou C, Katsambas AD. Origin, clinical presentation, and diagnoses of facial hypermelanoses. (8). *Dermatol Clin.* 2007;25:321