

PERFORATOR FLAPS FOR RECONSTRUCTION OF LOWER LIMB DEFECTS

Dr. Santosh Kumar

Senior Resident, Department of in-Charge Burn and Plastic Surgery, Government Medical College and Hospital, Bettiah, Bihar, India

Article Info: Received 23 September 2020; Accepted 28 October 2020

DOI: <https://doi.org/10.32553/ijmbs.v4i10.1516>

Corresponding author: Dr. Santosh Kumar

Conflict of interest: No conflict of interest.

Abstract

Background: Reconstruction of soft tissue defects in the lower third of the leg remains challenging. Anatomical constraints limit the local options available for complex defects especially lower third of leg. Local flaps based on perforator vessels are raising interest in reconstructive surgery of the limbs. We present our experience with perforator flaps for reconstruction of soft tissue defects in the lower limb.

Methods: This prospective cross-sectional study was carried out in the Department of Burn and Plastic Surgery at Government Medical College and Hospital, Bettiah, Bihar, India from Jan 2018 to August 2018. Total 20 patients with lower limb defects treated with various perforator flaps (both elective as well as emergency) were included in the study. A hand-held Ultrasound Doppler was used preoperatively and intra-operatively to detect the perforator vessels.

Results: Out of 20 patients, 6 patients in our study developed complications, out of which 3 patients had minor complications which included infection, wound dehiscence and congestion of flap. These patients did not require any secondary procedure and were managed conservatively. We witnessed partial flap loss in one and distal flap necrosis in two patients. In one of these three cases, the flap was repositioned to its native site, due to suspected colour changes intra-operatively and later after 3 days was successfully positioned on the desired site. The other two cases were managed with VAC and later STSG.

Conclusions: Perforator flaps may represent a good alternative to the free flaps in the areas where other local reconstructive procedures are not possible. This is a versatile technique and with decreased donor site morbidity limited to a single body area. There is a specific like to like soft tissue replacement leading to a better cosmetic and reconstructive outcome. The main drawback of the perforator flaps however is the higher risk of venous congestion.

Keywords: Perforator flaps, Propeller flaps, Lower limb reconstruction, Lower third leg reconstruction

Introduction:

Plastic surgery is a constant battle between blood supply and beauty. The end result of a reconstructive procedure is primarily attributable to the stability of the vascular component, which is fundamental in that it ensures survival and proper functioning of tissues that have been transferred to the recipient site.¹ The lower limb has always been known for poor wound healing and, since the first steps of the plastic surgery, as a scarce source of flaps for reconstruction. Soft tissue reconstruction of the lower limb is hence, challenging. Due to limited mobility and a paucity of overlying skin, even small soft tissue defects of the lower limb generally need flap coverage.² Before the introduction of microsurgery, surgeons had few reconstructive options such as local flaps (random skin flaps, muscular or musculocutaneous flaps) and performed cross-legs, immobilizing the limbs for weeks.³ A random

pattern flap has an indistinct perfusion pattern and is limited in size and mobility.⁴

Musculocutaneous flaps and muscle flaps with skin grafts such as from the gastrocnemius, soleus, and tibialis anterior can be used in the proximal and middle thirds of a pretibial defect.⁵ Unfortunately, the area least well served by these muscle flaps is the lower third of the leg. The fasciocutaneous flap reported by Ponten showed that long narrow flaps could be safely raised below the knee as long as the deep fascia was included.⁶ Ponten's flaps were not based on specific perforators and therefore could not be islanded. Free microvascular transfer is an answer to most of the difficult reconstructions but it is time consuming, requires microsurgical facility and expertise. After a long evolution of the reconstructive methods, the reappraisal of the works of Manchot and Salmon by Taylor and Palmer opened the era of perforator flaps. This era began in 1989, when Koshima and Soeda, and separately Kroll and

Rosenfield described the first applications of such flaps. Improvement in the anatomical knowledge on cutaneous, subcutaneous, and intramuscular vessels originating from major vascular axis of the limbs has allowed development of several types of perforator flaps, which today are commonly employed in clinical practice.⁷⁻⁹ With the development of perforator flaps newer and more reliable flaps have become available for lower limb reconstruction.¹⁰ According to the Gent consensus, perforator flaps are composed of skin and subcutaneous fat nourished by perforators arising from deep vascular systems, which reach the surface by passing mostly through muscle and intramuscular septa.¹¹ Although perforator flaps technique requires microsurgical dissection, it does not require vascular suturing and can thus be defined a microsurgical nonmicrovascular flap as reported by Georgescu et al.¹² Avoiding vascular sutures makes the surgical act quicker in comparison with microvascular flaps.

Material and Methods

This prospective cross-sectional study was carried out in the Department of Burn and Plastic Surgery at Government Medical College and Hospital, Bettiah, Bihar, India from Jan 2018 to August 2018, after taking the approval of the protocol review committee and institutional ethics committee.

Methodology

After taking informed consent detailed history was taken from the patient or the relatives if the patient was not in good condition. 20 patients with lower limb defects treated with various perforator flaps (both elective as well as emergency) during this period were included in the study.

The etiology, site, size and characteristics of the defect and surrounding area were analysed. In designing the flaps, the vascular axes and the distribution of the perforators which could sustain them were taken into consideration. A hand-held Ultrasound Doppler was used preoperatively and intra-operatively to detect perforator vessels in the donor site area. Perforator artery selection before flap harvesting was based on vessel size and distance to the area of the defect. Once the perforator was identified, the flap was designed around the perforator or perforators according to the location and size of the defect. The dimensions of the flap were based upon the size of defect and the movement

of the flap, taking into account the need to avoid excessive tension on the margins of the flap during suturing.

Surgical technique

The operations were performed using magnification loupes (3.5-4.0X) and microsurgical instruments. A tourniquet was inflated without prior exsanguinations. This maneuver facilitates identification of perforators as they remain filled with the blood. An exploratory incision along the margin of flap was made keeping the position of marked perforator in mind. The incision is made through the skin, subcutaneous tissue, deep fascia (sub-fascial approach) and the perforator vessel is directly visualized. The incision is initially always made from one side of the flap only to properly identify and assess the calibre of the perforator. If the perforator previously identified by Doppler is not adequate, we looked for another suitable perforator and the flap design was modified accordingly. Careful and meticulous dissection was done in a blunt way isolating the perforator. Usually the perforator had to be dissected for several centimeters to allow easier rotation or advancement. Pedicle traction during flap harvesting and positioning was avoided. Adequate release of all fascial strands around the perforator and dissection around the perforator in intermuscular or intramuscular plane to gain additional length were then carried out. This facilitates rotation of flap without kinking the perforator. After deflation of the tourniquet, hemostasis was performed and viability of flap was evaluated. Perfusion was checked before flap rotation by waiting a few minutes and irrigating with lukewarm saline solution in order to promote microcirculation recovery. The flap was then rotated on its perforator to varying degrees and inset into the defect after ensuring the viability and rechecking the vascularise while in desired position. Carefully positioned drains were then applied at the end of the procedure in some patients according to need. Drains were usually removed after 24 hrs. Bandaging was soft, to avoid compression over the flap, and the limb was held in an elevated position. A window was left uncovered for monitoring of colour and temperature without bandage removal. The donor sites/secondary defects in almost all the patients were grafted. Postoperatively the flaps were monitored. The parameters monitored included colour, temperature, margins, signs of poor perfusion/congestion, epidermal shrinking, blistering.

Results

Table 1:

Age	Sex	Etiology	Site of defect	Perforator flap source artery	Type/ Movement	No. of perforators	Size of flap	Duration of procedure (hrs)	Donor site Management	Complications
30	F	RTA	R. leg (lower third)	Peroneal	Propeller	1	9x6	2.3	STSG	nil
47	F	Exposed implant (post tumour resection)	L. leg (upper third)	Posterior tibial	Rotation	2	8X5	2.3	STSG	Nil
35	F	Operated S.C.C	L. gluteal region	Superior gluteal	Propeller	1	11x10	3	Primarily closed	congestion/ necrosis is of distal 2.4 cm
50	M	Unstable scar (post glass cut)	R. medial malleolus	Posterior tibial	Propeller	1	7x5	1.45	STSG	congestion/ one third necrosis
67	M	RTA	R. leg (upper third)	Anterior tibial	Propeller	1	5x5	1.25	STSG	Nil
68	M	RTA	L. leg (lower third)	Posterior Tibial	Rotation	1	9X7	2.3	STSG	Nil
52	M	Fall from tree (exposed implant)	L. leg (lower third)	Peroneal	Propeller	2	7x6	2.3	STSG	Infection
47	F	Cellulitis (Infection)	R. leg (upper third)	Posterior tibial	Transposition	1	8x5	2.15	STSG	Nil
62	F	RTA	L. leg (lower third)	Peroneal	Propeller	2	11x5	2.3	STSG	Nil
18	M	RTA	L. leg (middle third)	Anterior tibial	Propeller	1	10x4	1.45	STSG	Congestion
52	M	Fall	R. leg (upper third)	Posterior tibial	Propeller	1	10x5	3.15	STSG	Nil
57	M	Tin cut injury	L. leg (lower third)	Posterior tibial	Propeller	1	7x4	1.45	STSG	nil
19	M	Trophic ulcer EX	L. heel	Posterior tibial	Propeller	1	10x5	2.15	STSG	Nil
19	M	RTA	L. knee	Superior genicular	Rotation	1	18x12	3.40	STSG	congestion/necrosis of distal 2.4cm
58	F	RTA	L. medial malleolus	Posterior tibial	Propeller	1	8x4	1.45	STSG	Nil
7	M	Cellulitis	L. leg (middle third)	Anterior tibial	Advancement	2	7x4	2.15	STSG	Nil
34	M	Fall	R. leg (lower third)	Posterior tibial	Propeller	1	12x7	3.3	STSG	Nil
33	M	Osteomyelitis	L. leg (middle third)	Anterior tibial	Propeller	1	6x4	2.3	STSG	Nil
27	F	RTA	R. leg (lower third)	Anterior tibial	Transposition	1	10x9	2	STSG	Necrosis of distal 2.4cm
72	F	Burn	R. medial malleolus	Posterior tibial	Rotation	1	6x4	3	STSG	Nil

Discussion

The lower limb has always been known for poor wound healing and soft tissue reconstruction of the lower limb is challenging. The ideal reconstruction technique for both simple and complex defects of the lower limb should replace like to like tissue, minimize donor-site morbidity, preserve main vascular trunks, and reduce operating and hospitalization time. Perforator based flaps meet most of these requirements. The development of perforator flaps

in reconstructive microsurgery has been facilitated by improved knowledge of the arterial basis of flap perfusion. The subdermic vascular network is particularly rich and allows the harvesting of thin skin flaps. One single perforator vessel located in an eccentric position in relation to a skin paddle may support a large skin area thanks to the opening of potential vascular territories, which move to the peripheral border of the flap. The process of vascular adoption is promoted by the increase

of blood pressure, which occurs in the perforator artery after closure of subcutaneous and intramuscular branches during flap harvesting. One of the main characteristics of perforator flaps is their versatility, as the flap may be selected on the perforator artery according to defect type.

The present study was carried out on 20 patients who underwent reconstruction of lower limb defects with various perforator flaps. More than one-third of our patients had soft tissue loss following road traffic accident which has shown an increase in recent years and the usual mode was motor bike accident. The various etiologies in our study included road traffic accidents, fall from height, oncological resections, post infection/cellulitis debridement, firearm injury, comode injury, unstable scar, tin cut injury, trophic ulcer and burn. Road traffic accident (40%) was the commonest of all. Maximum cases in our study as well as those reported in literature were road traffic accidents, the reason being that there has been a tremendous increase in the number of vehicles and nowadays road traffic accidents are predicted to be the third leading contributor to the global burden of disease. The most common site of reconstruction in our study was the lower third of leg which constituted 35% of cases. The fact that lower third of leg is a difficult site for reconstruction with limited options, perforator flaps have been recommended by several clinical studies reported on the application of perforator based local flaps in lower limb reconstruction.¹³⁻¹⁷ The size of perforator flap that can be safely harvested has always been a point of argument and bone of contention for plastic surgeons all over the world. The length of the flap in our study ranged from 4 to 20 cm and the width from 3 to 11 cm. The maximum size of flap harvested in our study was 18×12 cm² for covering the knee joint in a young patient with an exposed knee joint following a road traffic accident. The large flap territory can be raised on a single perforator due to extensive axial communications between the perforators within the flap. Hyperperfusion in a perforator allows the capture of multiple adjacent perforasomes through direct and indirect lining vessels. Panse et al. in their study made an attempt to define the safe extent of local perforator flap for lower limb reconstruction by comparing it with the limb length of the patient and concluded that there is a six times more chance that a local perforator flap will necrosis if it is more than one-third of the limb length as compared to a flap which is less than one-third of the limb length.¹⁸ It is pertinent to mention here that there is still no standardization or reference for safe limit of a perforator flap.

Maximum flaps in our study were islanded and propelled into the defect as these flaps have the advantage of maximum gain in movement and increased arc of rotation. The operative time in most of studies ranges between 2 to

3 hours. Average duration of surgery in our study was 2.40 hours, with maximum duration of 3.40 hours and minimum duration of 1.25 hours. This makes the perforator flap reconstruction a preferred option for patients with comorbidities who might not be good candidates for longer duration surgeries. We witnessed partial flap loss in one and distal flap necrosis in two patients following postoperative venous congestion. In one of these three cases, the flap was repositioned to its native site, due to suspected colour changes and mild congestion intra-operatively and later after 72hours was successfully positioned on the desired site. The other two cases were managed with VAC and later STSG. The main reason for flap loss has been attributed to venous congestion which can occur due to kinking of the vein following flap rotation because of thinner wall as compared to that of the artery.

Conclusion

The perforator flaps in lower limb reconstruction are a viable option in the armamentarium of a reconstructive surgeon. This is a versatile technique and with decreased donor site morbidity limited to a single body area. These flaps do not involve sacrifice of any of the main arteries. They can cover very distal defects of the leg. There is a specific like to like soft tissue replacement leading to a better cosmetic and reconstructive outcome. The operative time taken for perforator flaps is not significantly higher than that for other fasciocutaneous flaps. Disadvantages of these flaps are that they have a limited role in larger defects, degloving injuries, and variable location of the perforators away from the soft tissue defect can sometimes be a hurdle. The main drawback of the perforator flaps however is the higher risk of venous congestion. Because the perforator venous wall is much thinner than the perforator arterial wall, it has a greater chance of venous congestion when rotated up to 180 degrees. The distal part of the flap which is used for defect coverage sometimes suffers from partial skin necrosis due to venous congestion. The problem of venous congestion can often be prevented with adequate dissection, loose suturing, postoperative dependent drainage and massage. Perforator flaps may represent a good alternative to the free flaps in the areas where other local reconstructive procedures are not possible. The favourable results reported in the literature, as well as the results of our personal experience for lower limb reconstruction, are encouraging. We believe that when the characteristics of the defect are suitable for treatment, this technique should be regarded as one of the possible and viable reconstructive option.

References

1. Gillies HD, Millard DR. The principles and art of plastic surgery. Boston: Little and Brown; 1957.

2. Godina M. Early microsurgical reconstruction of complex trauma of the extremities. *Plast Reconstr Surg.* 1986;78:285-92.
3. Jayes PH. Cross-leg flaps: a review of 60 cases. *Br J Plast Surg.* 1950;3:1-5.
4. Quaba O, Quaba A. Pedicled perforator flaps for the lower limb. *Semin Plast Surg.* 2006;20:103-11.
5. Fayman MS, Orak F, Hugo B, Berson SD. The distally based split soleus muscle flap. *Br J Plast Surg.* 1987;40:20-6.
6. Ponten B. The fasciocutaneous flap: its use in soft tissue defects of the lower leg. *Br J Plast Surg.* 1981;34:215-20.
7. Taylor GI. The angiosomes of the body and their supply to perforator flaps. *Clin Plast Surg.* 2003;30:331-42.
8. Koshima I, Moriguchi T, Ohta S. The vasculature and clinical application of the posterior tibial perforator-based flap. *Plast Reconstr Surg.* 1992;90:643-9.
9. Tanaka K, Matsumura H, Miyaki T. An anatomic study of the intermuscular septum of the lower leg; branches from the posterior tibial artery and potential for reconstruction of the lower leg and the heel. *J Plast Reconstr Aesthet Surg.* 2006;59:835-8
10. Hallock GG. Lower extremity muscle perforator flaps for lower extremity reconstruction. *Plast Reconstr Surg.* 2004;114:1123-30.
11. Schaverien M, Saint-Cyr M. Perforators of the lower leg: analysis of perforator locations and clinical application for pedicled perforator flaps. *Plast Reconstr Surg.* 2008;122:161-70.
12. Georgescu AV, Matei I, Ardelean F, Capota I. Microsurgical nonmicrovascular flaps in forearm and hand reconstruction. *Microsurgery.* 2007;27:1-7.
13. Acland RD, Schusterman M, Godina M, Eder E, Taylor GI, Carlisle I. The saphenous neurovascular free flap. *Plast Reconstr Surg.* 1981;67:763-74.
14. Wei FC, Mardini S. Free-style free flaps. *Plast Reconstr Surg.* 2004;114:910-6.
15. Taylor GI, Pan WR. Angiosomes of the leg: anatomic study and clinical implications. *Plast Reconstr Surg.* 1998;102:599-616.
16. Hyakusoku H, Yamamoto T, Fumiiri M. The propeller flap method. *Br J Plast Surg.* 1991;44:53-4.
17. Yildirim S, Taylan G, Akoz T. Freestyle perforatorbased V-Y advancement flap for reconstruction of soft tissue defects at various anatomical regions. *Ann Plast Surg.* 2007;58:501-6.
18. Panse NS, Bhatt YC, Tandale MS. What is safe limit of the perforator flap in lower extremity reconstruction? Do we have answers yet? *Plast Surg Int.* 2011; 2011:349-57