

CLINICAL PROFILE AND OUTCOME OF CHILDREN ADMITTED WITH STATUS EPILEPTICS

Dr. Sushil Kuamr Bakolia¹, Dr. Gajendra Kumar Verma², Dr. Dinesh Kumar Barolia³

¹MD (Paediatrics), Assistant Professor, Govt. S.K. Medical College & AG Hospital, Sikar (Rajasthan)

²MD (Paediatrics), Senior Resident, SMS Mc, Jaipur

³M Ch (Paediatric) Surgery, Assistant Professor, J L N Mc, Ajmer

Article Info: Received 05 August 2021; Accepted 19 September 2021

DOI: <https://doi.org/10.32553/ijmbs.v5i9.2207>

Corresponding author: Dr Dinesh Kumar Barolia

Conflict of interest: No conflict of interest.

Abstract

Background: Objective of this study was to study epidemiology and clinical profile of Status epilepticus

Methods: Hospital based cross sectional study conducted on 50 children All children aged between 1 month to 12 years who at presentation or during the PICU stay had convulsive status epilepticus - defined as continuous seizure activity or recurrent seizure activity without regaining consciousness lasting for >5 min.

Results: Among 50 children 56.00 % were in the age group were less than 5 years ,24 % were between 6-10 years ,20.00 % were above 10 yrs. The mean age group was 6.21±1.26 years. Incidence was higher in males (64%) when compared to females (36%). Generalized tonic clonic seizure were observed in 46 (92.00%) and partial seizure was noted in 4 (8%) of the children. About 36 (72%) of the children developed SE for the first time.

Conclusion: Status epilepticus is one of the common neurological emergency which requires admission to PICU. In our study epilepsy is one of the most common causes of status epilepticus. Early and appropriate treatment with anticonvulsants and use of mechanical ventilation may improve the outcome.

Keywords: Status epilepticus, mortality, clinical profile

Introduction

Status epilepticus (SE) is a common life-threatening emergency that requires prompt recognition and management.¹ SE can represent an exacerbation of a preexisting seizure disorder, the initial manifestation of a seizure disorder, or an insult other than a seizure disorder resulting in seizures.

The classical definition of SE requires that seizures (continuous or intermittent without return to baseline mental status) last for a minimum of 30 minutes. However, seizures which last longer than 5 minutes are unlikely to stop spontaneously.² As part of operational definition put forth within the past few years, status epilepticus is defined as continuous seizure activity or recurrent seizure activity without regaining consciousness, lasting for more than 5 minutes. The use of operational definition allows early treatment before the seizure becomes refractory to antiepileptic drugs.⁴ Prolonged status epilepticus can lead to various complications such as cardiac dysrhythmia, metabolic derangements, autonomic dysfunction, neurogenic pulmonary oedema, hyperthermia, pulmonary aspiration and even permanent neurological damage. Approximately 4-10% of children experience one episode of seizure within first 16 years of life. Approximately 30% of patients presenting with status epilepticus are having their first seizure. Status epilepticus is most common in children

younger than 5 years of age with an incidence of >100 per 100,000 children. Convulsive status epilepticus (CSE) is most common form of status epilepticus and accounts for about 90% of all SE in children. Mortality from status epilepticus varies from 3- 50% in different studies.^{3,4}

Material and Methods

Study Design:

Hospital based prospective study

Study Population:

All children aged between 1 month to 12 years who at presentation or during the PICU stay had convulsive status epilepticus - defined as continuous seizure activity or recurrent seizure activity without regaining consciousness lasting for >5 min.

Inclusion Criteria:

All children aged between 1 month to 12 years who at presentation or during the PICU stay had convulsive status epilepticus - defined as continuous seizure activity or recurrent seizure activity without regaining consciousness lasting for >5 min.

Exclusion Criteria:

Patients in whom the information regarding seizure duration will be incomplete or unclear.

Data Collection:

Informed consent will be obtained from parents or guardians of the children included in the study. The duration of status epilepticus will be ascertained from a reliable patient's relative or attendant, medical records and referring physician's note. After securing airway, breathing and circulation all the patients will be managed with standard treatment protocol.

Once the child will be stabilized, data which included age, sex, duration of seizures before and after admission, type and number of antiepileptic drugs (AEDs) used for control of status epilepticus (SE), history of previous seizure pattern, adherence to treatment, perinatal, developmental, family history and history of coexisting medical conditions will be recorded.

General physical examination and detailed neurological examination will be performed. Investigations like complete blood count, blood chemistries including serum calcium, random blood sugar, serum sodium, urea and creatinine, neuroimaging, CSF examination, Electroencephalography

(EEG) will be performed as required to ascertain etiology and guide management. Further, during the hospital stay, recurrence of seizures, subsequent need for intubation and mechanical ventilation and days spent in PICU will be noted. Type and etiology of status epilepticus will be classified as per report of the International League Against Epilepsy (ILAE) task force on classification of status epilepticus.

Refractory status epilepticus (RSE) was defined as seizures which persist despite the administration of two appropriate anticonvulsants at acceptable doses, with a minimum duration of status of 60 minutes (by history or on observation). Though febrile seizures are a part of acute symptomatic etiology, it has been considered separately for analysis as it is likely to erroneously amplify the severity of febrile seizures and dilute the severity of acute neurological insults.

Data Analysis:

All data were analyzed on EPI-info statistical software. Qualitative data were expressed in the form of proportion. Quantitative data were expressed in mean \pm SD. Qualitative data were compared by Chi square test. Unpaired t test will be used to infer the difference in means.

Results**Table 1: Demographic Profile**

		No of cases	Percentage
Age	0-5 Yrs	26	52.00
	6-10 Yrs	12	24.00
	>10 Yrs	12	24.00
Sex	Male	31	62.00
	Female	19	38.00

Among 50 children 52.00% were in the age group were less than 5 years, 24% were between 6-10 years, 24.00% were above 10 yrs. Incidence was higher in males (64%) when compared to females (36%).

Table 2: Clinical profile

Seizure type	GTCS	46	92.00
	Partial	4	8.00
Nausea / vomiting		9	18.00
Fever		28	56.00
Faecal and urinary incontinence		7	14.00
Headache		11	22.00
Developmental delay		13	26.00
Respiratory infection		5	10.00
Hospital stay		3.12 \pm 2.06 days	
Mortality		3	6.00

Generalized tonic clonic seizure were observed in 46 (92.00%) and partial seizure was noted in 4 (8%) of the children. About 36 (72%) of the children developed SE for the first time.

Discussion

In this study, it was found that children less than five years of age comprised the majority of the cases (56.00%). Other studies have also found a higher prevalence in the younger age group. As the first episode of convulsion has been theorized to be due to the underdeveloped mechanisms for control of seizure activity, there is a disruption of neuronal function with minimal abnormalities in younger children. Also, younger age is more vulnerable to acute etiologies including febrile seizures.^{5,6}

In present study, the sex distribution shows slightly higher incidence of status epilepticus in males than females. Although they were almost in equal proportions, many other studies done both in pediatrics and adults shows similar type of results Gulati S observed 22 (70%) patients out of 31 were male.⁷

It was observed that GTCS type of seizure is the commonest of all 4 types similar incidence was observed by Kwong KL⁸ and Gulati et al.⁷

Conclusion

Status epilepticus is one of the common neurological emergency which requires admission to PICU. In our study epilepsy is one of the most common causes of status

epilepticus. Early and appropriate treatment with anticonvulsants and use of mechanical ventilation may improve the outcome.

References

1. Pellock JM. Status epilepticus in children: update and review. *J Child Neurol.* 1994;9(2):27-35
2. Mikati MA, Hani AJ. Status epilepticus. In: Kliegman R, Behrman R, Nelson W. *Nelson textbook of pediatrics, 20e*; Elsevier; 2016:2854- 2855.
3. Trinka E, Cock H, Hesdorffer D, Rossetti AO, Scheffer IE, Shinnar S, et al. A definition and classification of status epilepticus-Report of the ILAE Task Force on Classification of Status Epilepticus. *Epilep.* 2015;56(10):1515-23.
4. Madhu PK, Krithika R. Convulsive status epilepticus in children: clinical profile and outcome in a tertiary care hospital. *Int J Contemp Pediatr* 2019;6:xxx-xx
5. Hauser WA. Status epilepticus: epidemiologic considerations. *Neurology.* 1990;40:S9-S13.
6. Treiman DM, Meyers PD, Walton NY, Collins JF, Colling C, Rowan AJ, Handforth A, Faught E, Calabrese VP, Uthman BM, Ramsay RE. A comparison of four treatments for generalized convulsive status epilepticus. *New Engl J Medicine.* 1998;339(12):792-8.
7. Gulati S, Kalra V, Sridhar MR. Status Epilepticus in Indian Children in a tertiary care center. *Indian J Pediatruics,* 2005;72:105-8
8. Kwong KL, Lee SL, Yung A, Wong VC. *J Child Health.* 1995(5):395-8.