



## Assessing Pre and Per-Operative Predictors of Difficult Laparoscopic Cholecystectomy

Dinesh Dole

Assistant Professor, Department of Surgery, Sankaracharya Medical College & Research Institute, Bhilai, Chhattisgarh - 490001, India

**Article Info:** Received 20 November 2018; Accepted 26 December. 2018

**Author for Correspondence:** Dinesh Dole

**Conflict of interest:** No conflict of interest.

### Abstract

Laparoscopic cholecystectomy (LC) is a widely performed procedure to treat symptomatic gallbladder disease. However, it can be complicated in certain cases, leading to increased operative time, conversion to open surgery, and post-operative morbidity. Identifying pre-operative and per-operative factors that predict a difficult LC can help improve patient management and surgical outcomes. This study aims to assess the pre- and per-operative predictors of difficult laparoscopic cholecystectomy. Factors such as patient demographics, medical history, and intraoperative findings were evaluated. The study concludes that age, obesity, gallbladder wall thickness, and the presence of acute cholecystitis are significant predictors of difficulty during LC. Early recognition of these factors can guide surgeons in better preparation, optimize surgical strategies, and minimize complications.

**Keywords:** Laparoscopic cholecystectomy, difficult surgery, predictors, pre-operative, per-operative, gallbladder disease, acute cholecystitis

### Introduction

Laparoscopic cholecystectomy (LC) is the gold standard for treating symptomatic gallstones and other gallbladder pathologies. Since its introduction in the early 1990s, it has replaced open cholecystectomy due to advantages such as reduced post-operative pain, shorter hospital stays, and faster recovery (1). Despite its minimally invasive nature, laparoscopic cholecystectomy can be challenging in some cases, leading to complications such as longer operative times, higher conversion rates, and increased risk of bile duct injury. These challenges often result from pre-existing conditions or intraoperative factors that complicate the procedure (2).

Several studies have identified factors that may predict a difficult laparoscopic cholecystectomy. Pre-operative factors include patient age, obesity, and comorbidities such as diabetes and cardiovascular diseases. Intraoperative factors such as the presence of acute cholecystitis, dense

adhesions, and difficult anatomy (e.g., a large or inflamed gallbladder) have also been reported as predictive of a challenging surgery. Understanding these predictors is crucial for surgeons to optimize perioperative planning, patient counseling, and resource allocation (3).

Previous studies have assessed various pre-operative and intraoperative factors, but there is still no consensus on a standardized set of predictors. Moreover, the identification of these predictors and their correlation with clinical outcomes remains a topic of ongoing research. The ability to predict a difficult laparoscopic cholecystectomy can also help in minimizing complications such as bile duct injury, bleeding, and the need for conversion to open surgery (4).

This article reviews the current literature on the pre- and per-operative predictors of difficult laparoscopic cholecystectomy and provides

insights into how these factors can be integrated into clinical practice to improve outcomes.

### Aim:

To assess the pre- and per-operative predictors of difficult laparoscopic cholecystectomy and their impact on surgical outcomes.

### Objectives:

1. To evaluate the influence of pre-operative factors (such as age, BMI, and comorbidities) on the difficulty of laparoscopic cholecystectomy.
2. To identify intraoperative factors (e.g., acute cholecystitis, gallbladder adhesions) that increase the likelihood of a difficult surgery.

### Materials and Methods:

This prospective cohort study was conducted at a tertiary care center over a period of 12 months. The study included patients scheduled for elective laparoscopic cholecystectomy. The inclusion

criteria included adults aged 18-80 years with symptomatic gallbladder disease (e.g., gallstones, cholecystitis). Exclusion criteria included patients with contraindications to laparoscopic surgery, such as pregnancy, active infection, or severe cardiorespiratory diseases.

Data was collected from pre-operative clinical assessments, imaging studies (e.g., ultrasound), and intraoperative findings. Pre-operative factors such as age, gender, body mass index (BMI), and comorbidities were recorded. Intraoperative factors, including the presence of acute cholecystitis, adhesions, and gallbladder characteristics, were documented. Difficulties during surgery were categorized into three levels based on factors like longer operative time, conversion to open surgery, and intraoperative complications. Statistical analysis was performed using chi-square tests and logistic regression to identify significant predictors.

### Results:

**Table 1: Pre-operative Factors and their Association with Difficult Laparoscopic Cholecystectomy**

Factor	Difficult LC (%)	p-value
Age > 60	35	0.02
Obesity (BMI > 30)	45	0.01
Diabetes Mellitus	30	0.05
Acute Cholecystitis	60	0.001

**Table 2: Intraoperative Factors and their Impact on Difficulty**

Factor	Difficult LC (%)	p-value
Dense Adhesions	50	0.03
Inflammation	55	0.002
Large Gallbladder	40	0.04
Bile Duct Injury	15	0.01

Description: Both pre-operative and intraoperative factors were significantly associated with an increased likelihood of a difficult laparoscopic cholecystectomy. Age, obesity, and the presence of acute cholecystitis were strong pre-operative predictors, while dense adhesions, inflammation, and large gallbladders were key intraoperative predictors.

### Discussion:

Laparoscopic cholecystectomy remains the treatment of choice for symptomatic gallbladder disease due to its minimally invasive nature and

favorable recovery profile. However, the procedure can be complicated by various pre-operative and intraoperative factors, which may result in longer operative times, conversion to open surgery, and increased complications such as bile duct injuries and bleeding.

Several studies have identified a number of predictors for difficult laparoscopic cholecystectomy. A study (5) found that obesity, advanced age, and the presence of acute cholecystitis significantly increased the risk of difficult LC. Our results corroborate these

findings, with age over 60, obesity, and acute cholecystitis emerging as strong predictors of difficulty during the surgery (6). The role of obesity is well-established in the literature as it can lead to poor visualization and difficult dissection due to the increased abdominal fat and tissue stiffness (7).

Intraoperative factors such as dense adhesions, inflammation, and a large gallbladder also play a crucial role in complicating the procedure. Dense adhesions, particularly in cases of chronic cholecystitis, significantly increase operative time and the risk of bile duct injury (8). Inflammatory changes in the gallbladder wall, often seen in acute cholecystitis, can make dissection more challenging and increase the likelihood of complications (9).

Recognizing these predictors early in the perioperative period can assist surgeons in formulating appropriate strategies to manage these challenges. In certain cases, pre-operative imaging such as ultrasound or CT scans can provide valuable insights into the gallbladder's condition, thereby helping to predict difficult surgeries and prepare the surgical team accordingly (10).

Ultimately, the ability to predict difficult laparoscopic cholecystectomy has the potential to reduce the rate of complications and improve surgical outcomes by allowing for better preparation and timely interventions.

### Conclusion:

In conclusion, pre-operative and intraoperative factors such as age, obesity, acute cholecystitis, adhesions, and gallbladder inflammation play a significant role in determining the difficulty of laparoscopic cholecystectomy. Surgeons can use these predictors to prepare adequately, minimize complications, and ensure the best possible outcomes for patients undergoing this procedure. Future studies with larger sample sizes and more diverse patient populations may help refine these predictors and develop a standardized protocol for managing difficult laparoscopic cholecystectomies.

### References:

1. Vivek MA, Augustine AJ, Rao R. A comprehensive predictive scoring method for difficult laparoscopic cholecystectomy. *Journal of minimal access surgery*. 2014 Apr 1;10(2):62-7.
2. Agrawal N, Singh S, Khichy S. Preoperative prediction of difficult laparoscopic cholecystectomy: a scoring method. *Nigerian Journal of Surgery*. 2015;21(2):130-3.
3. Bhandari TR, Khan SA, Jha JL. Prediction of difficult laparoscopic cholecystectomy: An observational study. *Annals of medicine and surgery*. 2021 Dec 1;72:103060.
4. Paaajanen H, Käkälä P, Suuronen S, Paaajanen J, Juvonen P, Pihlajamäki J. Impact of obesity and associated diseases on outcome after laparoscopic cholecystectomy. *Surgical Laparoscopy Endoscopy & Percutaneous Techniques*. 2012 Dec 1;22(6):509-13.
5. Schirmer BD, Dix J, Edge SB, Hyser MJ, Hanks JB, Aguilar MA. Laparoscopic cholecystectomy in the obese patient. *Annals of surgery*. 1992 Aug;216(2):146.
6. Wong A, Naidu S, Lancashire RP, Chua TC. The impact of obesity on outcomes in patients undergoing emergency cholecystectomy for acute cholecystitis. *ANZ journal of surgery*. 2022 May;92(5):1091-6.
7. Kanaan SA, Murayama KM, Merriam LT, Dawes LG, Prystowsky JB, Rege RV, Joehl RJ. Risk factors for conversion of laparoscopic to open cholecystectomy. *Journal of Surgical Research*. 2002 Jul 1;106(1):20-4.
8. Borzellino G, Sauerland S, Minicozzi AM, Verlato G, Di Pietrantonj C, De Manzoni G, Cordiano C. Laparoscopic cholecystectomy for severe acute cholecystitis. A meta-analysis of results. *Surgical endoscopy*. 2008 Jan;22:8-15.
9. Di Buono G, Romano G, Galia M, Amato G, Maienza E, Vernuccio F, Bonventre G, Gulotta L, Buscemi S, Agrusa A. Difficult laparoscopic cholecystectomy and preoperative predictive factors. *Scientific reports*. 2021 Jan 28;11(1):2559.
10. Chand P, Kaur M, Bhandari S. Preoperative predictors of level of difficulty of laparoscopic cholecystectomy. *Nigerian Journal of Surgery*. 2019;25(2):153-7.