"PREVALENCE OF ISOLATED HYPERTENSION AMONG ELDERLY PATIENTS ATTENDING PERIPHERAL HEALTH CENTRE OF TERTIARY CARE HOSPITAL OF INDIA."

Dr. Munish Kumar Sharma¹, Dr. Neeraj Gour², Dr. Sanjay Choudhary³, Dr. N K Goel⁴, Dr. Deepak Sharma⁵, Dr. Meenakshi Chaudhary⁶

¹Associate Professor, Dept of Community Medicine, GMCH sector-32, Chandigarh

²Associate Professor, Department of Community Medicine, SHKM Govt. Medical College, Mewat, Haryana, India

³Associate Professor, Department of Community Medicine, Govt. Medical College, Pali, Rajasthan, India

⁴Professor & Head, Dept of Community Medicine, GMCH sector-32, Chandigarh

⁵Assistant Professor, Dept of Community Medicine, GMCH sector-32, Chandigarh

⁶Post Graduate Resident, Dept of Pathology, Index Medical College & Hospital, Indore

Article Info: Received 22 November 2019; Accepted 16 December 2019

DOI: <https://doi.org/10.32553/ijmbs.v3i12.813>

Corresponding author: Dr. Sanjay Choudhary

Conflict of interest: No conflict of interest.

Abstract

Background: In this modern era, chronic non communicable diseases (NCDs) are a major global health challenge with numerous grave complications on the human body. Old age population tends to carry risk of almost all kind of diseases including non communicable diseases. Hypertension is one of biggest well known risk factor for many of non-communicable diseases usually found in the outdoor patients.

Method: This cross sectional study was undertaken to provide the data on the Prevalence & Pattern of hypertension amongst patients who were aged 60 years & above and who attended the Out Patients Department of the Rural Health & Training Centre of tertiary care hospital of Chandigarh city of India.

Results: Majority of patients belonging to age group 60-69 years were found to be diastolic hypertensive (24.46%). Majority of male patients were found to be diastolic hypertensive (28.43%) ($P < 0.05$). More or less same trend was found among female patients.

Conclusion: Hypertension was found to be highly prevalent among the outdoor elderly patients of Rural Health & Training Centre of Chandigarh, India. This study warrants the need of an early detection of hypertension, which can be accomplished by a periodic screening of the people.

Keywords: Hypertension; Elderly; India.

Introduction

In this modern era, chronic noncommunicable diseases (NCDs) are a major global health challenge with numerous grave complications on the human body. NCDs kill 40 million people each year, equivalent to 70% of all deaths globally. Each year, 15 million people die from a NCD between the ages of 30 and 69 years; over 80% of these premature deaths occur in low and middle income countries^[1]

Hypertension is the most common disorder which is being encountered in the outdoor patients^[2]. It is an established major risk factor and a leading cause of cardiovascular diseases worldwide. The prevalence of hypertension and its complications are increasing in the developing countries^[3]. Cardiovascular diseases

will be the largest cause of death and disability in India by 2020^[4]. The economic development, changes in the lifestyle and diet, and an increase in the life expectancy may be attributed to this rapid increase. Hypertension is an iceberg disease. The proportion of the awareness, treatment, and the control of hypertension are unacceptably low in the Indian adult population, as indicated from a plethora of studies^[5-8].

Thus, hypertension is emerging out as a major public health problem. The prevalence of hypertension has significantly increased more among urban communities than rural people. The prevalence of hypertension will increase even further, unless broad and effective preventive measures are implemented.

Prevention programmes and policies are in their infancy in India. Epidemiological studies are urgently needed to assess the prevalence of hypertension to determine the baseline, against which the future trends in the risk factor levels can be assessed and preventive strategies can be planned to promote health among the populations.^[9]

The present study was undertaken to provide the data on the prevalence of Isolated hypertension amongst patients who were aged 60 years and above and who attended the Out Patients Department of the Rural Health & Training Centre of tertiary care hospital of Chandigarh city of India.

Material and Methods:

Study Type: Observational secondary data based study

Study Duration: May 2018 to August 2018 covering a total of 245 (Male 102 & Female 143) patients aged 60 years & above.

Study Population: All the patients visited RHTC (Rural Health Training Centre) of GMCH (Govt. Medical College & Hospital), Chandigarh, India, aged 60 years & above.

Study Sample: A total of 245 patients were selected as sample size, who visited the RHTC during study period.

Participant Recruitment:

- Inclusion criteria: All the adults who were aged 60 years & above., who attended the Rural Health Training Centre of Govt. Medical College & Hospital, Chandigarh, India.
- Exclusion criteria: All the participants who were aged less than 60 years of age were excluded from the study. The patients who were non co-operative and those who refused to provide the necessary information were also excluded.

Data Collection:

This study was conducted between May 2018 and August 2018. A total of 245 patients (102 males, 143 females) gave their consent and participated in the study. A structured, pretested proforma was used to collect the data with regards to the socio-demographic characteristics and the blood pressure pattern. Blood pressure was measured by the trained staff in a bid to avoid any measurement bias. Calibrated BP instrument was used throughout the study period in a bid to curtail any instrumental bias.

Two measurements of the blood pressure of each study participant were made 30 minutes apart, with the participants in the sitting position, by using a mercury column sphygmomanometer. The blood pressure measurements were made on the subject's left arm by using a cuff of the appropriate size, at the level of the heart. The cuff pressure was inflated 30 mm Hg above the level at which the radial pulse disappeared and then it was deflated slowly at the rate of about 2mm per sec. In the cases where the two readings differed by over 10 mm of Hg, a third reading was obtained, and average of three measurements was taken. The pressures, at which the sound appeared and disappeared, were taken as the Systolic Blood Pressure (SBP) and the Diastolic Blood Pressure (DBP) respectively.

Operational Definitions:^[10]

- Persons in the age group of 18–59 years having systolic BP ≥ 140 mmHg and diastolic BP ≥ 90 mmHg or any level of BP in patients taking antihypertensive medication were considered as hypertensive.
- Persons in the age group of 60 or more than 60 years having systolic BP ≥ 150 mmHg and diastolic BP ≥ 90 mmHg or any level of BP in patients taking antihypertensive medication were considered as hypertensive.

Data processing and analysis:

Data collected were compiled and analyzed using suitable Statistical Software. The study was done using 95% confidence level. $P < 0.05$ was considered statistically significant. Pearson's Chi-square test was used to evaluate differences between groups for categorized variables.

Results:

This study was conducted on elderly Patients (Age 60 years & above) visited the Rural Health and training Centre of Govt. Medical College and Hospital Chandigarh. During the study period a total of 245 patients visited the health facility. Out of these all patients majority of patients were females (58.36%) and by far age distribution is concerned majority of patients (76.73%) were in the age group of 60 to 69 years. Association of age with systolic blood pressure was found to be non significant ($P > 0.05$). On the other hand association gender with systolic blood pressure was also found to be non significant ($P > 0.05$). (Table: 1)

Table 1: Showing Association of socio demographic characteristics with systolic blood pressure

Age group (in years)	Systolic Blood pressure (in mmHg)				Total N(%)
	<120 N (%)	120-139 N(%)	140-149 N(%)	>150 N(%)	
60-69	47(25)	87(46.27)	32(17)	22(11.7)	188 (76.73)
70 & above	19(33.33)	21(36.84)	05(8.77)	12(21.05)	57 (23.26)
Total	66(26.93)	108(44.08)	37(15.10)	34(13.87)	245 (100)
Chi square:6.74 & P value: 0.0807 (> 0.05)					
Gender					
Male	32(31.37)	43(42.15)	14(13.72)	13(12.74)	102(41.63)
Female	34(23.77)	65(45.45)	23(16.08)	21(8.57)	143 (58.36)
Total	66(26.93)	108(44.08)	37(15.10)	34(13.87)	245 (100)
Chi square: 1.80 & P value: 0.614 (> 0.05)					

Study has also tried to find out Association of diastolic blood pressure with age and gender. No significant association was found out of diastolic blood pressure with both age ($P > 0.05$) and Gender respectively ($P > 0.05$). (Table: 2)

Table 2: Showing Association of socio demographic characteristics with diastolic blood pressure

Age group (in years)	Diastolic Blood pressure (in mmHg)				Total N (%)
	<80 N (%)	89-89 N (%)	90-99 N (%)	>100 N (%)	
60-69	74(39.36)	68(36.17)	38(20.21)	08(4.25)	188 (76.73)
70 & above	22(38.59)	19(33.33)	10(17.54)	06(10.52)	57 (23.26)
Total	96(39.18)	87(35.51)	48(19.59)	14(5.71)	245 (100)
Chi square:3.27 & P value: 0.351 (> 0.05)					
Gender					
Male	35(34.31)	38(37.25)	20(19.60)	09(8.82)	102(41.63)
Female	61(42.65)	49(34.26)	28(19.58)	05(3.49)	143 (58.36)
Total	96(39.18)	87(60.83)	48(19.59)	14(5.71)	245(100)
Chi square:4.16 & P value: 0.24 (> 0.05)					

By far prevalence of hypertension among age groups is concerned majority of patients belonging to age group 60-69 years were found to be diastolic hypertensive (24.46%) ($P < 0.05$). Whereas more or less equal distribution of prevalence was found in 70 & above age groups for systolic & diastolic hypertension ($P < 0.05$). Overall diastolic hypertension prevalence was more than systolic hypertension ($P > 0.05$). (Table: 3)

Table 3: Age group wise distribution of Systolic & Diastolic hypertension

Age group (in years)	Systolic Hypertension	Diastolic Hypertension	Total	Chi square	P value
60-69	22 (11.7)	46(24.46)	188 (46.73)	10.33	0.0013
70 & above	12(21.05)	16(28.07)	57 (23.26)	0.76	0.383
Total	34(13.87)	62 (25.30)	245 (100)	10.16	0.0014

By far prevalence of hypertension among gender is concerned majority of male patients were found to be diastolic hypertensive (28.43%) ($P < 0.05$). More of less same trend was found among female patients. (Table: 4)

Table 4: Gender wise distribution of Systolic & Diastolic hypertension.

Gender	Systolic Hypertension	Diastolic Hypertension	Total	Chi square	P value
Male	13(12.7)	29(28.43)	102(41.63)	7.72	0.0054
Female	21(14.68)	33(23.07)	143(58.36)	3.29	0.698
Total	34(13.8)	62(25.30)	245(100)	10.30	0.0013

Discussion:

The prevalence of hypertension ranged from 5-15% between 1960-1990^[11]. It had increased to 20-36% in the past decade^[12]. The factors which were attributable to this rising trend were the rapid urbanization, lifestyle changes, dietary changes and the increased life expectancy. The high prevalence of hypertension (44.46%) in the current study, confirmed this increasing trend. A compatible prevalence of hypertension was reported (46%) in a study which was conducted among the outdoor patients of an urban health center of Kolkata^[13].

This observational study was conducted on elderly Patients (Age 60 years & above) visited the rural health and training center of Government Medical College and Hospital Chandigarh. During the study period a total of 245 patients visited the health facility.

This study has found out that age and gender was not significantly associated with systolic blood pressure. More or less same kind of observation was also noticed with diastolic blood pressure where age and gender were not significantly associated.

Prakash J et al also found significant association of hypertension with the age group in a Rohtak based study.^[14]

Similarly, statistically significant association of hypertension with age group was observed in studies by Abu-Aisha et al.^[15], Ganesh et al.^[16], Ramakrishnan et al.^[17], Dhungana et al.^[18] among general population of age group 18–70 years in Kathmandu, Nagammanavar et al.^[19] among bank employees of Bellary city, Karnataka, and Ismail et al.^[20].

The proportion of hypertension, as well as the mean systolic and diastolic blood pressures, were found to increase steadily with an increase in age. These findings were coherent with those which had been reported in the studies which were conducted among the urban adults of Mumbai^[21] and Lucknow^[22]. Such changes of blood pressure with age, might be due to the changes in the vascular system. Cross-sectional surveys, as well as prospective observational cohort studies, have consistently demonstrated a positive relation between the age and blood pressure in most of the populations, with diverse geographical, cultural and socioeconomic characteristics^[23].

Present study has found out that hypertension (both systolic & Diastolic variant) was more prevalent

among age group of 60 to 69 years. More or less same kind of observation was also noticed in a study conducted by Mehmood SE et al.^[9]

Diastolic hypertension was found to be more prevalent among elderly age groups ($P < 0.05$). A study conducted at Rohtak by Prakash J et al^[14] & study conducted by Mehmood SE et al^[9] at Lucknow has found age as an independent significant predictor of hypertension.

Prevalence of diastolic hypertension among male patients was found to be almost double as compared to systolic hypertension ($P < 0.05$). Almost similar observations was found among female patients ($P > 0.05$). Study conducted at Lucknow^[9] has found prevalence of hypertension more among females contrary to our study. They have also found that gender is not a significant predictor of hypertension.

Comparable observations (48.4% in females and 47.5% in males) were reported among the urban adults of Mumbai^[21]. A female preponderance was also observed by Kabir et al., among the outdoor patients of Dhaka Medical College^[24]. In contrast, a greater proportion of hypertension was observed among males (42.9%) as compared to females (34.2%), among the urban adults of Lucknow, India^[22].

Hospital based study, single day measurement of blood pressure without taking history of other potential confounders were some of the limitation of this study.

Conclusion:

Hypertension was found to be highly prevalent among the outdoor patients of Rural Health & Training Centre of Chandigarh. This study warrants the need of an early detection of hypertension, which can be accomplished by a periodic screening of the people, regularly at the hospital as well as at the community levels.

References:

1. World Health Organization. Non Communicable Disease Fact Sheet. Geneva: World Health Organization. Available from: <http://www.who.int/mediacentre/factsheets/fs355/en/>. [Last accessed on 2019 Sep 04].
2. Cherry DK, Woodwill DA. National ambulatory medical care survey: 2000 summary. *Adv Data*. 2002; 328: 1-32.
3. The World Health Report 1999: The double burden: emerging epidemics and persistent problems.

- Geneva: WHO; 1999. Available from: <http://www.who.org/>, accessed on June 2, 2009.
4. Gupta R. Trends in hypertension epidemiology in India. *Journal of Human Hypertension*. 2004; 18:73–78.
 5. Quasem I, Shetye MS, Alex SC, Nag AK, Sarma PS, Thankappan KR, et al . Hypertension Study Group. Prevalence, awareness, treatment and control of hypertension among elderly in Bangladesh and India: a multicentric study. *Bulletin of the World Health Organization* 2001;79(6):490-500.
 6. Bharucha NE, Kuruvilla T. Hypertension in the Parsi community of Bombay: A study on prevalence, awareness and compliance to treatment. *BMC Public Health*. 2003;3:1-5.
 7. Deshmukh PR, Gupta SS, Bharambe MS, Maliye C, Kaur S, Garg BS. Prevalence of hypertension, its correlates and levels of awareness in rural Wardha, Central India. *Journal of Health & Population in Developing Countries*. 2005;1-12.
 8. Mohan V, Deepa M, Farooq S, Datta M, Deepa R. Prevalence, awareness and control of hypertension in Chennai – The Chennai Urban Rural Epidemiology Study (CURES – 52). *JAPI*. 2007;55: 326-32.
 9. Mahmood SE, Prakash D, Srivastava JP, Zaidi ZH, Bhardwaj P. Prevalence of hypertension amongst adult patients attending out patient department of Urban Health Training Centre, Department of Community Medicine, Era's Lucknow Medical College and Hospital, Lucknow. *Journal of clinical and diagnostic research: JCDR*. 2013 Apr;7(4):652.
 10. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 Evidence-based guideline for the management of high blood pressure in adults: Report from the panel members appointed to the eighth joint national committee (JNC 8). *JAMA* 2014;311:507-20.
 11. Gupta R. Meta-analysis of prevalence of hypertension in India. *Indian Heart J*. 1997; 49: 43-48.
 12. Mohan V, Deepa M, Farooq S, Datta M, Deepa R. Prevalence, Awareness and Control of Hypertension in Chennai - The Chennai Urban Rural Epidemiology Study (CURES – 52). *JAPI*. 2007; 55: 326-32.
 13. Soumya Deb, Aparajita Dasgupta. A Study on Risk Factors of Cardiovascular Diseases in an Urban Health Center of Kolkata. *Indian Journal of Community Medicine*. 2008; 33 (4): 275-75.
 14. Parkash J, Kalhan M, Singhanian K, Punia A, Kumar B, Kaushal P. Prevalence of hypertension and its determinants among policemen in a City of Haryana, India. *Int J App Basic Med Res* 2019;9:143-7.
 15. Abu-Aisha H, Elhassan EA, Khamis AH, Abu-Elmaali A. Hypertension and obesity in police forces households in Khartoum, Sudan: A pilot report – Part of the “police forces hypertension, diabetes, renal insufficiency, and thyroid derangements (HyDRIT) study”, Sudan. *Sudan J Public Health* 2008;3:17-25.
 16. Ganesh KS, Naresh AG, Bammigatti C. Prevalence and risk factors of hypertension among male police personnel in urban Puducherry, India. *Kathmandu Univ Med J (KUMJ)* 2014;12:242-6.
 17. Ramakrishnan J, Majgi SM, Premarajan KC, Lakshminarayanan S, Thangaraj S, Chinnakali P. High prevalence of cardiovascular risk factors among policemen in Puducherry, South India. *J Cardiovasc Dis Res* 2013;4:112-5.
 18. Dhungana RR, Pandey AR, Bista B, Joshi S, Devkota S. Prevalence and associated factors of hypertension: A Community-based cross-sectional study in municipalities of Kathmandu, Nepal. *Int J Hypertens* 2016;2016:1656938.
 19. Nagammanavar R, Somashekhar G, Reddy CS, Kumar P, Raghavendra B. A study of prevalence and risk factors of hypertension among the bank employees of Bellary city: A cross-sectional study. *J Sci* 2015;5:459-66.
 20. Ismail IM, Kulkarni AG, Kamble SV, Borker SA, Rekha R, Amruth M. Prevalence of hypertension and its risk factors among bank employees of Sullia Taluk, Karnataka. *Sahel Med J* 2013;16:139-43.
 21. Gupta PC, Gupta R, Pednekar MS. Hypertension prevalence and blood pressure trends in 88 653 subjects in Mumbai, India. *Journal of Human Hypertension*. 2004; 18:907–10.
 22. Yadav S, Boddula R, Genitta G, Bhatia V, Bansal B, Kongara S. Prevalence and risk factors of pre-hypertension & hypertension in an affluent north Indian population. *Indian J Med Res*. 2008; 128: 712-20.
 23. Hypertension control. Technical Report Series: World Health Organization. 1996. Report No.: 862.
 24. A Kabir, TK Barman, NA Yousuf, N Fatima, J Banik. Prevalence of Hypertension among the Patients who attend medicine outdoor of Dhaka Medical College Hospital. *J Medicine*. 2007; 8 : 49-52