CONTEMPORARY TREATMENT OF ALK POSITIVE NON SMALL CELL LUNG CANCER PATIENT WITH BRAIN METASTASES: OUR EXPERIENCE

Crvenkova S¹, Popova M¹

¹ University Clinic of Radiotherapy and Oncology, Faculty of Medicine, Skopje

Article Info: Received 21 December 2019; Accepted 10 January, 2020

DOI: <https://doi.org/10.32553/ijmbs.v4i1.835>

Corresponding author: Crvenkova Simonida

Conflict of interest: No conflict of interest.

Abstract

Purpose: The aim of this study is to show alectinib efficacy and safety, with focus on alectinib intracranial efficacy.

Case presentation: We report on a 46-year-old woman diagnosed as non small cell lung cancer harboring echinoderm microtubule-associated protein-like 4-anaplastic lymphoma kinase fusion gene. She has intracranial metastases and poor performance status of 3. We treated this patients with alectinib after whole brain irradiation and discontinuation of chemotherapy after two 2 cycles due to progressive brain and liver metastases. Good response was obtained with improving of her performance status without adverse events.

Conclusions: We recommend alectinib as a treatment approach for ALK+ NSCLC patients, especially those with CNS metastases at the time of the diagnosis and poor PS.

Key words: CNS metastasis, ALK+ NSCLC patients, alectinib

Introduction

Lung cancer remains a worldwide epidemic. Approximately 1.2 million people die from lung cancer each year (1). NSCLC represents over 80% of all lung cancers and 60% to 70% of the patients with NSCLC suffer from stage III or IV disease. The molecular profile of the tumor currently determines the therapeutic strategy for advanced lung cancer. Distinctive chromosomal rearrangements in the ALK gene (ALK-positive) were first described in 2007 and occur in approximately 2%–5% of patients with nsclc (2,3). The most common ALK rearrangement is a fusion between the N-terminal half of *eml4* and the intracellular kinase domain of *alk* (*eml4-alk*) (4,5), leading to an active oncogenic driver. Other variations of ALK rearrangements exist. Additional ALK-related oncogenic drivers include point mutations in the kinase domain and *alk* overexpression (6,7). Patients with ALK-positive nsclc are typically younger and tend to be light or never-smokers (8). Incidence of brain metastases (BMs) is higher in patients with ALK-positive NSCLC and among those patients up to 50-60% will develop BMs during the course of their disease (8). Alectinib is potent second-generation ALK inhibitor and was shown to be effective for a broad spectrum of ALK rearrangement and ALK mutations. The aim of this study is to show alectinib efficacy and safety with focus on alectinib intracranial efficacy.

Case presentation

A 46-year-old woman with light smoking history (15 years 10 cigarettes per day) was admitted to our hospital because of cerebral symptomatology, dizziness and headache for at list 15 days. Patient has a poor performans status (PS) of 3. Chest X-ray and computed tomography (CT) scans showed a tumor lesion in the inferior left lung lobe. Bronchoscopy established a pathological diagnosis of adenocarcinoma with a component of signet ring cells. Along with CT scan of abdomen and brain magnetic resonance imaging (MRI) scan, reveled multiple liver and brain metastases. The patient was finally diagnosed as having stage IV lung adenocarcinoma cTNM=T3N2M1b (LIV, BRA). Patient was first treated with whole brain irradiation (WBI) to a 30 Gy with 300 cGy daily dose and two cycles of first-line chemotherapy with carboplatine/gemcitabine. Although chemotherapy yielded temporary tumor regression, progressive brain and liver metastases led to discontinuation of chemotherapy after the two courses. In turn alectinib 300 mg bid was administered, because of the positive results of EML-4-ALK fusion protein and gene from immunohistochemical analysis. Alectinib is covered by the national health insurance system in our country. She is currently responding to alectinib 600 mg daily. After 9 months, CT and MRI showed lung and brain disappearance and substantial tumor regression in liver (Fig.1, 2, 3) The patient continues to be treated with alectinib without critical adverse events. At the latest follow-up, 10 months after commencing alectinib treatment, there was no evidence of progression or any remarkable toxicity.

Figure1: Computed tomography scans depicting the tumor response at the inferior left lung lobe, 9 months after alectinib treatment (A pretreatment lung foci, B positive response to the alectinib after 9 months)

Figure 2: Magnetic resonance imaging scans depicting the response of intracranial metastases after alectinib treatment (A pretreatment metastatic foci, B positive response to the alectinib after 9 months)

Figure 3: Computed tomography scans depicting the tumor response at the liver, 9 months after alectinib treatment (A pretreatment liver metastatic foci, B positive response to the alectinib after 9 months)

Discussion

Our study describe the case of an ALK-positive NSCLC patient with sever cerebral simptomatology and poor PS. There has been little clinical benefit of WBI and chemotherapy in NSCLC with poor PS and BMs with a median OS of about 3 months (9) The indication for molecular targeted therapy for NSCLC patients harboring corresponding target genes therefore needs to be determined separately from that for cytotoxic chemotherapy for non-selected population. In the context ALK inhibitor developed and has demonstrated a systemic efficacy and strongly improved out-comes in patients with ALK-positive advanced NSCLC in comparison with chemotherapy. First line median progression-free survival PFS was longer with crizotinib in comparison with chemotherapy (10.9 versus 7 months). However, the intracranial efficacy of crizotinib is poor, due to poor blood-brain barrier (BBB) penetration (10,11). There was a need for development of other ALK inhibitors to improve intracranial disease control and enlarge the spectrum of ALK mutations targeted. For these reasons, the second generation ALK inhibitors ceritinib, alectinib and brigatinib and the third-generation ALK inhibitor lorlatinib were developed (12). More recently, Kodama and colleagues showed alectinib to have a higher antitumor activity than crizotinib in intracranial tumors in mouse model of EML4-ALK-positive NSCLC due a higher BBB penetration (13). Alectinib is not P-glycoprotein substrate and this may play

a role in the higher BBB penetration. P-glycoprotein overexpression has indeed been showed to be a mechanism of resistance to ALK inhibitors, especially in the brain (14). Alectinib efficacy in patients with BMs was also assessed in phase III clinical trials. In the ALUR study (15), 24 patients in the alectinib arm and 16 patients in the chemotherapy arm had baseline measurable CNS metastases. The icORR was significantly higher with alectinib 54.2% versus chemotherapy 0% $p < 0.001$. Several real-life retrospective case series confirmed the high efficacy of alectinib on BMs and leptomeningeal metastases LM (16,17). Moreover, Ou and colleagues reported the cases of the two patients with ALK-positive NSCLC with BMs who received stereotactic radiosurgery (SRS) to the brain prior to alectinib treatment. Both patients had radiation necrosis presenting as pseudo-progression confirmed by neurosurgery and pathologic examination (18). This specific brain evolution after SRS and alectinib has to be known to avoid incorrect classification into progressive disease and alectinib discontinuation. Therefore we chose alectinib for our patient. At present, crizotinib and alectinib are available as ALK inhibitors in our country. Results from approach in J-ALEX and ALEX trials provided further evidence of alectinib's systemic and CNS efficacy, with complete CNS response rates of 38% in patients with measurable CNS lesions at baseline. In the ALEX ITT population, the cumulative incidence rate (CIR) of CNS progression,

considering the challenging risks of non-CNS progression and death, was 9.4% with alectinib versus 41.4% with crizotinib (19,20) Our patients with ALK-positive NSCLC who was previously treated with cytotoxic drugs and radiotherapy and still was without response with alectinib treatment demonstrated a good response regarding brain metastasis. At the latest follow-up, 10 months after commencing alectinib treatment, there was no evidence of progression or any remarkable toxicity.

Conclusion

Considering its efficacy and tolerability in this case, we recommend alectinib as a best treatment approach for ALK+ NSCLC patients, especially those with CNS metastasis at the time of the diagnosis and poor PS.

Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images.

References:

1. WHO Statistics. <http://www.who.int/mediacentre/factsheets/fs297/en/>) accessed 21 December 2017)
2. Salomon B, Varella-Garcia M and Camidge DR. ALK gene rearrangements a new therapeutic target in a molecularly defined subset of non-small cell lung cancer. *J Thorac Oncol* 2009;4: 1450-1454
3. Soda M, Choi YL, Enomoto M, et al. Identification of the transforming EML4-ALK fusion gene in NSCLC. *Nature* 2007;448:561-566
4. Rikova K, Guo A, Zeng Q, et al. Global survey of phosphotyrosine signaling identifies oncogenic kinases in lung cancer. *Cell* 2007;131:1190-1203
5. Kris M, Johnson BE, Berry LD, et al. Using multiplexed assays of oncogenic drivers in lung cancers to select targeted drugs. *JAMA* 2014;311:1998-2006
6. Shaw AT, Yeap BY, Mino-Kenudson M, et al. Clinical features and outcome of patients with NSCLC who harbor EMLA-4-ALK. *J Clin Oncol* 2009;27:4247-4253
7. Kwak EL, Bang YJ, Camidge DR, et al. Anaplastic lymphoma kinase inhibition in NSCLC. *N Engl J Med* 2010;363:1693-1703
8. Passaro K, Lazzari C, Karachaliou N et al. Personalized treatment in advanced ALK-positive NSCLC; from bench to clinical practice. *Onco Targets Ther.* 2016; 9:6361-6376
9. Rossi A, Alectinib for ALK-positive NSCLC. *Exp Rev Clin Pharmacol* 2016;9:1005-1013
10. Costa DR, Kobayashi S, Pandya SS, et al. CSF concentration of anaplastic lymphoma kinase inhibitor crizotinib. *J Clin Oncol* 2011;29:c443-c445.
11. Metro G, Lunardi G, Floridi P, et al. CSF concentration of crizotinib in two ALK-positive NSCLC patients with CNS metastases deriving clinical benefit from treatment. *J Thorac Oncol* 2015;10:c26-c27.
12. Ricciuti B, De Gioglio A, Mecca c, et al. Precision medicine against ALK positive NSCLC: beyond crizotinib. *Med Oncol* 2018;35:72
13. Kodama T, Hasegawa M, Takanashi K, et al. Antitumor activity of selective ALK inhibitor alectinib in models of intracranial metastasis. *Cancer Chemother Pharmacol* 2014;74:1023-1028
14. Katayama R, Sakashita T, Yanagitani N, et al. P-glycoprotein mediates ceritinib resistance in anaplastic lymphoma kinase-rearranged NSCLC. *EBio Medicine* 2016;3:54-66
15. Novello S, Mazieres J, Oh JJ, et al. Alectinib versus chemotherapy in crizotinib-pretreated anaplastic lymphoma kinase (ALK)-positive NSCLC: results from the phase III ALUR study. *Ann Oncol* 2018;29: 1409-1416.
16. Metro G, Lunardi G, Bennati C, et al. Alectinib's activity against CNS metastases from ALK-positive NSCLC a single institution case series. *J Neurooncol* 2016;129 355-361
17. Gainor JF, Sherman CA, Willoughby K, et al. Alectinib salvages CNS relapses in ALK-positive lung cancer patients previously treated with crizotinib and ceritinib. *J Thorac Oncol* 2015;10 232-236
18. Ou SH, Klemperer SJ, Azada MC, et al. Radiation necrosis presenting as pseudoprogression (PsP) during alectinib treatment of previously irradiated brain metastases in ALK-positive NSCLC implications for disease assessment and management Lung cancer 2015;88:355-359
19. Tran PN and Klemperer SJ. Focus on alectinib and competitor compounds for second line therapy in ALK-rearranged NSCLC. *Front Med (Lausanne)* 2016; 3:65